Principal stage Germination			Zadok's Code	
Barley can germinate at 34 degrees F 0 Dry kernel 0.1 Start of imbibition (water absorption) 0.5 Radicle emerged 0.7 Coleoptile emerged 0.9 Leaf just at coleoptile tip Seedling Development 0.1 First leaf at least 50 percent emerged 0.2 (12 days after emergence) 0.3 Third leaf at least 50 percent emerged 0.4 Fourth leaf at least 50 percent emerged 0.5 Fifth leaf at least 50 percent emerged 0.6 Fifth leaf at least 50 percent emerged 0.7 First leaf through coleoptile 0.8 Fifth leaf at least 50 percent emerged 0.9 Main shoot plus one tiller visible 0.1 Main shoot plus one tillers 0.2 Main shoot plus one tillers 0.3 Main shoot plus two tillers 0.4 Main shoot plus four tillers 0.5 Main shoot plus four tillers 10.5 Main shoot plus five tillers 10.5 Main shoot plus four tillers 10.6 Main shoot plus four tillers 10.7 First node detectable 10.8 Second node detectable 10.9 Fig leaf just visible 10.9 Fig leaf collar just visible 10.9 Fig leaf collar just visible 10.1 Figs leaf sheath extending 10.2 Boot swollen 10.3 Boot just beginning to swell (45 days) 10.4 First awns visible 10.5 Boot swollen 10.7 Fig leaf sheath opening 10.9 First awns visible 10.1 First spikelet of head just visible 10.2 Beginning off lowering 10.3 Beginning off lowering 10.4 First spikelet of head emerged 10.5 Half of head emerged 10.6 Half of florets have flowered 10.9 Flowering complete 10.1 Kernel watery ripe 10.2 Early milk 10.3 Early milk 10.4 Early dough 10.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 10.7 Hard dough, head losing green color (Aviv) 10.9 Approximate physiological maturity 10.1 Kernel hard (difficult to divide with fingernail) 10.1 Thumbnail cannot dent kernel, harvest ripe (84-90)	-	-	Description	
Barley can germinate at 34 degrees F O Dry kernel** 0.1 Start of imbibition (water absorption) 0.5 Radicle emerged 0.7 Coleoptile emerged 0.9 Leaf just at coleoptile tip **Seedling Development** O First leaf through coleoptile 0.1 First leaf at least 50 percent emerged Second leaf at least 50 percent emerged (12 days after emergence) 0.3 Third leaf at least 50 percent emerged (12 days after emergence) 0.4 Fourth leaf at least 50 percent emerged 0.5 Fifth leaf at least 50 percent emerged 10.5 Fifth leaf at least 50 percent emerged 11 Wain shoot only 12 O Main shoot plus one tiller visible 13 O Main shoot plus two tillers 14 O Main shoot plus two tillers 15 O Main shoot plus three tillers (16-23 days) 16 O Main shoot plus five tillers 17 O Main shoot plus five tillers 18 O Main shoot plus five tillers 19 O Main shoot plus five tillers 10 O Main shoot plus four tillers 10 O Main shoot p	stage	stage	•	
0 Dry kernel 0.1 Start of imbibition (water absorption) 0.5 Radicle emerged 0.7 Coleoptile emerged 0.9 Leaf just at coleoptile tip Seedling Development 0 First leaf through coleoptile 0.1 First leaf at least 50 percent emerged 0.2 Second leaf at least 50 percent emerged 12 days after emergence) 0.3 Third leaf at least 50 percent emerged 0.4 Fourth leaf at least 50 percent emerged 0.5 Fifth leaf at least 50 percent emerged 0.6 First leaf at least 50 percent emerged 0.7 First leaf at least 50 percent emerged 0.8 Main shoot only 0 Main shoot only 0 Main shoot plus two tillers 0.2 Main shoot plus two tillers 0.3 Main shoot plus four tillers 0.4 Main shoot plus four tillers 0.5 Main shoot plus free tillers (16-23 days) 0.1 First node detectable 0.2 Second node detectable 0.2 Second node detectable 0.2 Second node detectable 0.9 Flag leaf just visible 0.9 Flag leaf collar just visible (38-39 days) 8 DOT Cycle 0.1 Flag leaf sheath extending 0.3 Boot swollen 0.7 Flag leaf sheath extending 0.9 First awns visible Head Emergence (58 days) 0.1 First spikelet of head just visible 0.1 Beginning of flowering 0.1 Beginning of flowering 0.1 Beginning of flowering 0.1 Beginning of flowering 0.1 Reginning flowered 0.1 Reginning flowering 0.1 Reginning flowering 0.1 Reginning of flowering 0.1 Seginning of flowering 0.2 Half of florets have flowered 0.3 Barly milk 0.5 Medium milk 0.7 Late milk 0.6 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity 0.1 Kernel hard (difficult to divide with fingernail) 0.1 Kernel hard (difficult to divide with fingernail)				
0.1 Start of imbibition (water absorption) 0.5 Radicle emerged 0.7 Coleoptile emerged 0.9 Leaf just at coleoptile tip Seedling Development 0 First leaf through coleoptile 0.1 First leaf at least 50 percent emerged 0.2 Second leaf at least 50 percent emerged 0.3 Third leaf at least 50 percent emerged 0.4 Fourth leaf at least 50 percent emerged 0.5 Fifth leaf at least 50 percent emerged 0.5 Fifth leaf at least 50 percent emerged 0.6 Fourth leaf at least 50 percent emerged 0.7 First leaf at least 50 percent emerged 0.8 First leaf at least 50 percent emerged 0.9 Main shoot plus two tillers 0.1 Main shoot plus one tiller visible 0.2 Main shoot plus two tillers 0.3 Main shoot plus three tillers (16-23 days) 0.4 Main shoot plus five tillers 0.5 Main shoot plus five tillers 0.5 Main shoot plus five tillers 0.6 Main shoot plus five tillers 0.7 Fiag leaf just visible 0.9 Flag leaf onde detectable 0.1 Fiag leaf just visible (38-39 days) 0.1 First node detectable 0.2 Second node detectable 0.3 Boot cycle 1 Fiag leaf sheath extending 0.3 Boot just beginning to swell (45 days) 0.5 Boot swollen 0.7 Flag leaf sheath opening 0.9 First awns visible 0.9 First awns visible 0.9 First awns visible 0.9 First awns visible 0.0 First spikelet of head emerged 0.7 Three-fourths of head emerged 0.7 Three-fourths of head emerged 0.7 Three-fourths of head emerged 0.9 Head emergence complete 0.9 Head emergence complete 0.9 Head emergence complete 0.9 Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.2 Beding of flowering 0.3 Early milk 0.5 Medium milk 0.7 Late milk 0.6 Medium milk 0.7 Late milk 0.8 First placel depending on weather) 0.9 Approximate physiological maturity 0.9 Approximate physiological maturity 0.1 Mernel hard (difficult to divide with fingernail) 0.1 Thumbnail cannot dent kernel, harvest ripe (84-90)				
0.5 Radicle emerged 0.7 Coleoptile emerged 0.9 Leaf just at coleoptile tip Seedling Development 0 First leaf through coleoptile 0.1 First leaf at least 50 percent emerged 0.2 Second leaf at least 50 percent emerged 0.3 Third leaf at least 50 percent emerged 0.4 Fourth leaf at least 50 percent emerged 0.5 Fifth leaf at least 50 percent emerged 0.5 Fifth leaf at least 50 percent emerged 0.6 Main shoot only 0.1 Main shoot plus one tiller visible 0.2 Main shoot plus two tillers 0.3 Main shoot plus three tillers (16-23 days) 0.4 Main shoot plus four tillers 0.5 Main shoot plus fure tillers Stem Elongation (16-31 days) 0.1 First node detectable 0.2 Second node detectable 0.2 Second node detectable 0.7 Flag leaf just visible 0.9 Flag leaf collar just visible (38-39 days) 80ot Cycle 0.1 Flag leaf sheath extending 0.3 Boot just beginning to swell (45 days) 0.9 Flag leaf sheath extending 0.9 First awns visible 0.7 Flag leaf sheath opening 0.9 First spikelet of head just visible 0.7 Flag leaf sheath opening 0.9 First spikelet of head just visible 0.3 One-fourth of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.2 Beginning of flowering 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 4 Air field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 0.1 Kernel hard (difficult to divide with fingernail) 0.2 Thumbnail cannot dent kernel, harvest ripe (84-90	0		,	
0.7 Coleoptile emerged 0.9 Leaf just at coleoptile tip Seedling Development 0.1 First leaf through coleoptile 0.1 First leaf at least 50 percent emerged 0.2 Second leaf at least 50 percent emerged (12 days after emergence) 0.3 Third leaf at least 50 percent emerged 0.4 Fourth leaf at least 50 percent emerged 0.5 Fifth leaf at least 50 percent emerged 0.6 Fifth leaf at least 50 percent emerged 0.7 Fifth leaf at least 50 percent emerged 0.8 Main shoot only 0.1 Main shoot plus to percent emerged 0.2 Main shoot plus to percent emerged 0.3 Main shoot plus to percent emerged 0.4 Main shoot plus to percent emerged 0.5 Main shoot plus to percent emerged 0.6 Main shoot plus to percent emerged 0.7 Main shoot plus to percent emerged 0.8 Main shoot plus to percent emerged 0.9 First node detectable 0.1 First node detectable 0.2 Second node detectable 0.2 Second node detectable 0.3 Third node detectable 0.9 Flag leaf collar just visible (38-39 days) 800 Event Second Percent Second Seco				
0.9 Leaf just at coleoptile tip Seedling Development 0 First leaf through coleoptile 0.1 First leaf at least 50 percent emerged 0.2 Second leaf at least 50 percent emerged (12 days after emergence) 0.3 Third leaf at least 50 percent emerged 0.4 Fourth leaf at least 50 percent emerged 0.5 Fifth leaf at least 50 percent emerged 0.5 Fifth leaf at least 50 percent emerged 0.1 Main shoot plus four tillers 10 Main shoot plus one tiller visible 10.2 Main shoot plus two tillers 10.3 Main shoot plus three tillers (16-23 days) 10.4 Main shoot plus five tillers 10.5 Main shoot plus five tillers 10.6 Main shoot plus five tillers 10.7 First node detectable 10.8 Second node detectable 10.9 Flag leaf just visible 10.9 Flag leaf collar just visible (38-39 days) 10.1 First node detectable 10.2 Second node detectable 10.3 Boot just beginning to swell (45 days) 10.4 Flag leaf sheath extending 10.5 Boot swollen 10.7 Flag leaf sheath extending 10.9 First awns visible 10.7 Flag leaf sheath opening 10.9 First spikelet of head just visible 10.1 First spikelet of head just visible 10.3 One-fourth of head emerged 10.4 Head Emergence (58 days) 10.5 Half of florets have flowered 10.9 Head emergence complete Flowering (Not readily visible in barley) 10.1 Reginning of flowering 10.1 Reginning of flowering 10.2 Half of florets have flowered 10.3 Early milk 10.5 Medium milk 10.6 Set arly dough 10.7 Hard dough, head losing green color (Aviv) 10.9 Approximate physiological maturity 10.1 Kernel hard (difficult to divide with fingernail) 10.1 Kernel hard (difficult to divide with fingernail) 10.1 Kernel hard (difficult to divide with fingernail)				
1 First leaf through coleoptile 0.1 First leaf at least 50 percent emerged 0.2 Second leaf at least 50 percent emerged (12 days after emergence) 0.3 Third leaf at least 50 percent emerged 0.4 Fourth leaf at least 50 percent emerged 0.5 Fifth leaf at least 50 percent emerged 10.5 Fifth leaf at least 50 percent emerged 10.6 Main shoot only 10.1 Main shoot plus one tiller visible 10.2 Main shoot plus two tillers 10.3 Main shoot plus three tillers (16-23 days) 10.4 Main shoot plus four tillers 10.5 Main shoot plus four tillers 10.5 Main shoot plus four tillers 10.6 Main shoot plus four tillers 10.7 First node detectable 10.8 Second node detectable (16-31 days) 10.9 Fiag leaf just visible 10.9 Fiag leaf sheath extending 10.3 Boot just beginning to swell (45 days) 10.1 Fiag leaf sheath extending 10.3 Boot just beginning to swell (45 days) 10.5 Boot swollen 10.7 Fiag leaf sheath opening 10.9 First awns visible 10.1 First spikelet of head just visible 10.3 One-fourth of head emerged 10.5 Half of head emerged 10.6 Half of head emerged 10.7 Three-fourths of head emerged 10.9 Head emergence complete 10.1 Beginning of flowering 10.1 Beginning of flowering 10.1 Beginning of flowering 10.1 Kernel watery ripe 10.1 Kernel watery ripe 10.2 Kernel watery ripe 10.3 Early milk 10.5 Medium milk 10.5 Medium milk 10.7 Late milk 10.8 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 10.1 Kernel hard (difficult to divide with fingernail) 10.1 Kernel hard (difficult to divide with fingernail) 10.2 Thumbnail cannot dent kernel, harvest ripe (84-90)		0.9		
0.1 First leaf at least 50 percent emerged				
1 Second leaf at least 50 percent emerged (12 days after emergence) 0.3 Third leaf at least 50 percent emerged 0.4 Fourth leaf at least 50 percent emerged 0.5 Fifth leaf at least 50 percent emerged 10.5 Fifth leaf at least 50 percent emerged 10.6 Main shoot plus percent emerged 10.1 Main shoot plus one tiller visible 10.2 Main shoot plus two tillers 10.3 Main shoot plus four tillers 10.5 Main shoot plus four tillers 10.5 Main shoot plus five tillers 10.5 Main shoot plus four tillers 10.6 Main shoot plus four tillers 10.7 Flag leaf just visible 10.9 Flag leaf just visible 10.9 Flag leaf collar just visible (38-39 days) 10.1 First node detectable 10.2 Second node detectable 10.3 Boot cycle 10.1 Flag leaf sheath extending 10.3 Boot just beginning to swell (45 days) 10.5 Boot swollen 10.7 Flag leaf sheath opening 10.9 First awns visible 10.3 One-fourth of head emerged 10.4 First spikelet of head just visible 10.3 One-fourth of head emerged 10.5 Half of head emerged 10.7 Three-fourths of head emerged 10.9 Head emergence complete 10.1 Beginning of flowering 10.2 Beginning of flowering 10.3 Early milk 10.5 Medium milk 10.7 Late milk 10.5 Medium milk 10.7 Late milk 10.8 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 10.1 Mernel hard (difficult to divide with fingernail) 10.1 Kernel hard (difficult to divide with fingernail) 10.2 Thumbnail cannot dent kernel, harvest ripe (84-90)	1	0		
1		0.1		
1 (12 days after emergence) 0.3 Third leaf at least 50 percent emerged 0.4 Fourth leaf at least 50 percent emerged 0.5 Fifth leaf at least 50 percent emerged 10 Main shoot only 0.1 Main shoot plus one tiller visible 0.2 Main shoot plus two tillers 0.3 Main shoot plus four tillers 0.4 Main shoot plus five tillers 0.5 Main shoot plus five tillers 10.5 Main shoot plus five tillers 10.6 Stem Elongation (16-31 days) 10.1 First node detectable 10.2 Second node detectable (16-31 days) 10.3 Third node detectable (16-31 days) 10.4 Flag leaf just visible 10.7 Flag leaf just visible 10.9 Flag leaf collar just visible (38-39 days) 10.1 Flag leaf sheath extending 10.3 Boot just beginning to swell (45 days) 10.5 Boot swollen 10.7 Flag leaf sheath opening 10.9 First awns visible 10.1 First spikelet of head just visible 10.3 One-fourth of head emerged 10.4 First spikelet of head just visible 10.5 Half of head emerged 10.7 Three-fourths of head emerged 10.9 Head emergence complete 10.1 Beginning of flowering 10.1 Beginning of flowering 10.2 Kernel watery ripe 10.3 Early milk 10.5 Medium milk 10.7 Late milk 10.8 Dough Development in Kernel 10.9 Flowering (20 days maximum to very hard kernel in the field depending on weather) 10.1 Hard dough, head losing green color (Aviv) 10.9 Approximate physiological maturity 10.1 Kernel hard (difficult to divide with fingernail) 10.1 Kernel hard (difficult to divide with fingernail) 10.1 Kernel hard (difficult to divide with fingernail)		0.2	, -	
0.4 Fourth leaf at least 50 percent emerged 0.5 Fifth leaf at least 50 percent emerged Tillering (15 days) 0 Main shoot only 0.1 Main shoot plus one tiller visible 0.2 Main shoot plus two tillers 0.3 Main shoot plus three tillers (16-23 days) 0.4 Main shoot plus five tillers 0.5 Main shoot plus five tillers 0.5 Main shoot plus five tillers 0.6 Main shoot plus five tillers 0.7 Flag leaf (16-31 days) 0.1 First node detectable 0.2 Second node detectable 0.7 Flag leaf just visible 0.9 Flag leaf collar just visible (38-39 days) 8 Boot Cycle 1.1 Flag leaf sheath extending 0.3 Boot just beginning to swell (45 days) 0.5 Boot swollen 0.7 Flag leaf sheath opening 0.9 First awns visible Head Emergence (58 days) 1.1 First spikelet of head just visible 0.3 One-fourth of head emerged 0.5 Half of head emerged 0.7 Three-fourths of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 1.1 Beginning of flowering 1.2 Beginning complete 1.3 Beginning complete 1.4 Milk Development in Kernel 1.5 Medium milk 0.7 Late milk 1.6 Dough Development in Kernel 1.7 Alard dough, head losing green color (Aviv) 1.8 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 1.2 Control of the field depending on weather) 1.3 Early dough 1.4 Kernel hard (difficult to divide with fingernail) 1.5 Kernel hard (difficult to divide with fingernail) 1.6 Kernel hard (difficult to divide with fingernail) 1.7 Thumbnail cannot dent kernel, harvest ripe (84-90)				
Tillering (15 days) O Main shoot only 0.1 Main shoot plus one tiller visible 0.2 Main shoot plus two tillers 0.3 Main shoot plus three tillers (16-23 days) 0.4 Main shoot plus four tillers 0.5 Main shoot plus four tillers 5 Main shoot plus four tillers 5 Main shoot plus four tillers 5 Main shoot plus five tillers 6 Main shoot plus twisible 6 Main shoot plus twisible 7 Main shoot plus twisible 8 Main shoot plus twisible 8 Main shoot plus twisible 9 Main shoot plus twisible 10 Main shoot plus twisible (16-31 days) 10 Main shot tillers 10 Main shot twillers 10 Main shot tillers 10 Main shot tillers 10 Main shot till				
Tillering (15 days) O Main shoot only O.1 Main shoot plus one tiller visible O.2 Main shoot plus two tillers O.3 Main shoot plus three tillers (16-23 days) O.4 Main shoot plus four tillers O.5 Main shoot plus five tillers Stem Elongation (16-31 days) O.1 First node detectable O.2 Second node detectable (16-31 days) O.3 Third node detectable (16-31 days) O.7 Flag leaf just visible O.9 Flag leaf collar just visible (38-39 days) Boot Cycle O.1 Flag leaf sheath extending O.3 Boot just beginning to swell (45 days) O.5 Boot swollen O.7 Flag leaf sheath opening O.9 First awns visible Head Emergence (58 days) O.1 First spikelet of head just visible O.3 One-fourth of head emerged O.7 Three-fourths of head emerged O.9 Head emergence complete Flowering (Not readily visible in barley) O.1 Beginning of flowering O.1 Beginning of flowering Alalf of florets have flowered O.9 Flowering complete Milk Development in Kernel O.1 Kernel watery ripe O.3 Early milk O.5 Medium milk O.7 Late milk Dough Development in Kernel O.8 Soft dough (20 days maximum to very hard kernel in the field depending on weather) O.7 Hard dough, head losing green color (Aviv) O.9 Approximate physiological maturity Ripening O.1 Kernel hard (difficult to divide with fingernail) Thumbnail cannot dent kernel, harvest ripe (84-90)				
2 Main shoot only 0.1 Main shoot plus one tiller visible 0.2 Main shoot plus two tillers 0.3 Main shoot plus four tillers 0.5 Main shoot plus four tillers 0.5 Main shoot plus five tillers 0.6 Main shoot plus five tillers 0.7 First node detectable 0.8 Second node detectable 0.9 Flag leaf just visible 0.9 Flag leaf just visible 0.9 Flag leaf collar just visible (38-39 days) 800 Cycle 0.1 Flag leaf sheath extending 0.3 Boot just beginning to swell (45 days) 0.5 Boot swollen 0.7 Flag leaf sheath opening 0.9 First awns visible Head Emergence (58 days) 0.1 First spikelet of head just visible 0.3 One-fourth of head emerged 0.7 Three-fourths of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.9 Flowering complete Milk Development in Kernel 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk 0.7 Late milk 0.8 Soft dough (20 days moximum to very hard kernel in the field depending on weather) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 0.1 Kernel hard (difficult to divide with fingernail) 0.2 Thumbnail cannot dent kernel, harvest ripe (84-90)				
2		0		
2			,	
0.3 Main shoot plus three tillers (16-23 days) 0.4 Main shoot plus five tillers 0.5 Main shoot plus five tillers Stem Elongation (16-31 days) 0.1 First node detectable 0.2 Second node detectable (16-31 days) 0.3 Third node detectable 0.7 Flag leaf just visible 0.9 Flag leaf collar just visible (38-39 days) 800 Cycle 0.1 Flag leaf sheath extending 0.3 Boot just beginning to swell (45 days) 0.5 Boot swollen 0.7 Flag leaf sheath opening 0.9 First awns visible Head Emergence (58 days) 10.1 First spikelet of head just visible 0.3 One-fourth of head emerged 0.7 Three-fourths of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.2 Half of florets have flowered 0.9 Flowering complete Milk Development in Kernel 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 8 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) Thumbnail cannot dent kernel, harvest ripe (84-90			·	
0.4 Main shoot plus four tillers 0.5 Main shoot plus five tillers Stem Elongation (16-31 days) 0.1 First node detectable 0.2 Second node detectable 0.7 Flag leaf just visible 0.9 Flag leaf collar just visible (38-39 days) Boot Cycle 0.1 Flag leaf sheath extending 0.3 Boot just beginning to swell (45 days) 0.5 Boot swollen 0.7 Flag leaf sheath opening 0.9 First awns visible Head Emergence (58 days) 0.1 First spikelet of head just visible 0.3 One-fourth of head emerged 0.5 Half of head emerged 0.7 Three-fourths of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.1 Beginning of flowered 0.9 Flowering complete Milk Development in Kernel Kernel watery ripe 0.1 Kernel watery ripe 0.3 Early milk 0.7 Late milk Dough Development in Kernel 8 O.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) Thumbnail cannot dent kernel, harvest ripe (84-90)	2			
0.5 Main shoot plus five tillers Stem Elongation (16-31 days) 0.1 First node detectable 0.2 Second node detectable 0.3 Third node detectable 0.7 Flag leaf just visible 0.9 Flag leaf collar just visible (38-39 days) Boot Cycle 0.1 Flag leaf sheath extending 0.3 Boot just beginning to swell (45 days) 0.5 Boot swollen 0.7 Flag leaf sheath opening 0.9 First awns visible Head Emergence (58 days) 0.1 First spikelet of head just visible 0.3 One-fourth of head emerged 0.7 Three-fourths of head emerged 0.7 Three-fourths of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.5 Half of florets have flowered 0.9 Flowering complete Milk Development in Kernel 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 8 O.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) Flowering cannot dent kernel, harvest ripe (84-90)				
Stem Elongation (16-31 days) 0.1 First node detectable 0.2 Second node detectable (16-31 days) 0.3 Third node detectable 0.7 Flag leaf just visible 0.9 Flag leaf collar just visible (38-39 days) Boot Cycle 0.1 Flag leaf sheath extending 0.3 Boot just beginning to swell (45 days) 0.5 Boot swollen 0.7 Flag leaf sheath opening 0.9 First awns visible Head Emergence (58 days) 0.1 First spikelet of head just visible 0.3 One-fourth of head emerged 0.5 Half of head emerged 0.7 Three-fourths of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.5 Half of florets have flowered 0.9 Flowering complete Milk Development in Kernel 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 8 O.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 7 Kernel hard (difficult to divide with fingernail) 7 Thumbnail cannot dent kernel, harvest ripe (84-90)			· ·	
3 O.1 First node detectable O.2 Second node detectable (16-31 days) O.3 Third node detectable O.7 Flag leaf just visible O.9 Flag leaf collar just visible (38-39 days) Boot Cycle O.1 Flag leaf sheath extending O.3 Boot just beginning to swell (45 days) O.5 Boot swollen O.7 Flag leaf sheath opening O.9 First awns visible Head Emergence (58 days) O.1 First spikelet of head just visible O.3 One-fourth of head emerged O.5 Half of head emerged O.7 Three-fourths of head emerged O.9 Head emergence complete Flowering (Not readily visible in barley) O.1 Beginning of flowering O.9 Flowering complete Flowering complete Milk Development in Kernel O.1 Kernel watery ripe O.3 Early milk O.5 Medium milk O.7 Late milk Dough Development in Kernel O.3 Early dough O.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) O.7 Hard dough, head losing green color (Aviv) O.9 Approximate physiological maturity Ripening O.1 Kernel hard (difficult to divide with fingernail) Thumbnail cannot dent kernel, harvest ripe (84-90)				
3	3	0.1		
0.7 Flag leaf just visible 0.9 Flag leaf collar just visible (38-39 days) Boot Cycle 0.1 Flag leaf sheath extending 0.3 Boot just beginning to swell (45 days) 0.5 Boot swollen 0.7 Flag leaf sheath opening 0.9 First awns visible Head Emergence (58 days) 0.1 First spikelet of head just visible 0.3 One-fourth of head emerged 0.5 Half of head emerged 0.7 Three-fourths of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.5 Half of florets have flowered 0.9 Flowering complete Milk Development in Kernel 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 8 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 7 Thumbnail cannot dent kernel, harvest ripe (84-90)		0.2	Second node detectable (16-31 days)	
0.9 Flag leaf collar just visible (38-39 days) Boot Cycle 0.1 Flag leaf sheath extending 0.3 Boot just beginning to swell (45 days) 0.5 Boot swollen 0.7 Flag leaf sheath opening 0.9 First awns visible Head Emergence (58 days) 0.1 First spikelet of head just visible 0.3 One-fourth of head emerged 0.5 Half of head emerged 0.7 Three-fourths of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.5 Half of florets have flowered 0.9 Flowering complete Milk Development in Kernel 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 8 O.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 7 Thumbnail cannot dent kernel, harvest ripe (84-90)		0.3	Third node detectable	
## Boot Cycle O.1 Flag leaf sheath extending		0.7		
4 0.1 Flag leaf sheath extending 0.3 Boot just beginning to swell (45 days) 0.5 Boot swollen 0.7 Flag leaf sheath opening 0.9 First awns visible Head Emergence (58 days) 0.1 First spikelet of head just visible 0.3 One-fourth of head emerged 0.5 Half of head emerged 0.7 Three-fourths of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.5 Half of florets have flowered 0.9 Flowering complete Milk Development in Kernel 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 8 O.3 Early dough Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) Thumbnail cannot dent kernel, harvest ripe (84-90)		0.9		
4 0.3 Boot just beginning to swell (45 days) 0.5 Boot swollen 0.7 Flag leaf sheath opening 0.9 First awns visible Head Emergence (58 days) 0.1 First spikelet of head just visible 0.3 One-fourth of head emerged 0.5 Half of head emerged 0.7 Three-fourths of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.5 Half of florets have flowered 0.9 Flowering complete Milk Development in Kernel 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 8 O.3 Early dough 0.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 7 Thumbnail cannot dent kernel, harvest ripe (84-90)				
4 0.5 Boot swollen 0.7 Flag leaf sheath opening 0.9 First awns visible Head Emergence (58 days) 0.1 First spikelet of head just visible 0.3 One-fourth of head emerged 0.5 Half of head emerged 0.7 Three-fourths of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.5 Half of florets have flowered 0.9 Flowering complete Milk Development in Kernel 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 8 O.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 Thumbnail cannot dent kernel, harvest ripe (84-90)	4			
0.7 Flag leaf sheath opening 0.9 First awns visible Head Emergence (58 days) 0.1 First spikelet of head just visible 0.3 One-fourth of head emerged 0.5 Half of head emerged 0.7 Three-fourths of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.5 Half of florets have flowered 0.9 Flowering complete Milk Development in Kernel 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 8 O.5 Early dough 0.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 Thumbnail cannot dent kernel, harvest ripe (84-90)				
O.9 First awns visible Head Emergence (58 days)				
Head Emergence (58 days) 0.1 First spikelet of head just visible 0.3 One-fourth of head emerged 0.5 Half of head emerged 0.7 Three-fourths of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.5 Half of florets have flowered 0.9 Flowering complete Milk Development in Kernel Kernel watery ripe 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 8 O.3 Early dough 0.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 Thumbnail cannot dent kernel, harvest ripe (84-90)				
5 D.1 First spikelet of head just visible 0.3 One-fourth of head emerged 0.5 Half of head emerged 0.7 Three-fourths of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.5 Half of florets have flowered 0.9 Flowering complete Milk Development in Kernel 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 0.3 Early dough 0.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 Thumbnail cannot dent kernel, harvest ripe (84-90)		0.5		
5 O.3 One-fourth of head emerged O.5 Half of head emerged O.7 Three-fourths of head emerged O.9 Head emergence complete Flowering (Not readily visible in barley) O.1 Beginning of flowering O.5 Half of florets have flowered O.9 Flowering complete Milk Development in Kernel O.1 Kernel watery ripe O.3 Early milk O.5 Medium milk O.7 Late milk Dough Development in Kernel O.3 Early dough O.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) O.7 Hard dough, head losing green color (Aviv) O.9 Approximate physiological maturity Ripening O.1 Kernel hard (difficult to divide with fingernail) O.2 Thumbnail cannot dent kernel, harvest ripe (84-90)		0.1		
5 0.5 Half of head emerged 0.7 Three-fourths of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 0.1 Beginning of flowering 6 0.5 Half of florets have flowered 0.9 Flowering complete Milk Development in Kernel 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 0.3 Early dough 0.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 Thumbnail cannot dent kernel, harvest ripe (84-90)				
0.7 Three-fourths of head emerged 0.9 Head emergence complete Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.5 Half of florets have flowered 0.9 Flowering complete Milk Development in Kernel 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 8 0.3 Early dough 0.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 0.2 Thumbnail cannot dent kernel, harvest ripe (84-90)	5	0.5	-	
Flowering (Not readily visible in barley) 0.1 Beginning of flowering 0.5 Half of florets have flowered 0.9 Flowering complete Milk Development in Kernel Comparison of the standard process of th		0.7		
6 0.1 Beginning of flowering 0.5 Half of florets have flowered 0.9 Flowering complete Milk Development in Kernel O.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 8 0.3 Early dough Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 0.2 Thumbnail cannot dent kernel, harvest ripe (84-90)				
6 0.5 Half of florets have flowered 0.9 Flowering complete Milk Development in Kernel 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 8 0.3 Early dough Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 0.2 Thumbnail cannot dent kernel, harvest ripe (84-90)				
7 Plowering complete Milk Development in Kernel 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 8 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 Thumbnail cannot dent kernel, harvest ripe (84-90)	6			
7				
7 0.1 Kernel watery ripe 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel Early dough Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 0.2 Thumbnail cannot dent kernel, harvest ripe (84-90)		0.9		
7 0.3 Early milk 0.5 Medium milk 0.7 Late milk Dough Development in Kernel 8 0.3 Early dough 0.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 0.2 Thumbnail cannot dent kernel, harvest ripe (84-90)				
8 O.5 Medium milk 0.7 Late milk Dough Development in Kernel Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 10.2 Thumbnail cannot dent kernel, harvest ripe (84-90)	7		, ·	
8 Dough Development in Kernel 0.3 Early dough Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 0.2 Thumbnail cannot dent kernel, harvest ripe (84-90)			,	
8 Dough Development in Kernel 0.3 Early dough Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 0.2 Thumbnail cannot dent kernel, harvest ripe (84-90)				
8 0.3 Early dough 0.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 0.2 Thumbnail cannot dent kernel, harvest ripe (84-90)				
8 0.5 Soft dough (20 days maximum to very hard kernel in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 0.2 Thumbnail cannot dent kernel, harvest ripe (84-90)				
8 in the field depending on weather) 0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 Thumbnail cannot dent kernel, harvest ripe (84-90)				
0.7 Hard dough, head losing green color (Aviv) 0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 7 Thumbnail cannot dent kernel, harvest ripe (84-90)	8	0.5		
0.9 Approximate physiological maturity Ripening 0.1 Kernel hard (difficult to divide with fingernail) 9 Thumbnail cannot dent kernel, harvest ripe (84-90)		0.7		
Properties 8 Properties 9 Prope				
9 0.1 Kernel hard (difficult to divide with fingernail) Thumbnail cannot dent kernel, harvest ripe (84-90				
U.Z		0.1	Kernel hard (difficult to divide with fingernail)	
days)	9	0.2	Thumbnail cannot dent kernel, harvest ripe (84-90	
This is a normal harley cycle which is dependent on the Meteorological			• •	

This is a normal barley cycle which is dependent on the Meteorological conditions throughout its life cycle. When inspecting at the end of the 12th Biblical lunar month we look for quantities in various locations that are at the 8.3-8.5 stages.